

HOW MANY IDEALS WHOSE QUOTIENT RINGS ARE GORENSTEIN EXIST?

NAOKI ENDO

ABSTRACT. For an Ulrich ideal in a Gorenstein local ring, the quotient ring is again Gorenstein. Aiming to further develop the theory of Ulrich ideals, this paper investigates a naive question of how many non-principal ideals whose quotient rings are Gorenstein exist in a given Gorenstein ring. The main result provides that the number of such graded ideals in a symmetric numerical semigroup ring R coincides with the conductor of the semigroup. We furthermore provide a complete list of non-principal graded ideals I in R whose quotient rings R/I are Gorenstein.

1. INTRODUCTION

We investigate a question of how many non-principal ideals whose residue class rings are Gorenstein exist in Gorenstein rings. The motivation of this query comes from the desire to develop the study of *Ulrich* ideals. The notion of Ulrich ideals is one of the modifications of that of *stable* maximal ideals introduced in 1971 by his monumental paper [13] of J. Lipman. The present one was formulated by S. Goto, K. Ozeki, R. Takahashi, K.-i. Watanabe, and K.-i. Yoshida [5] in 2014, where the authors developed and consolidated the basic theory of Ulrich ideals. For example, if A is a Gorenstein local ring and that for Ulrich ideals I and J , the equality $I = J$ holds, provided $\mathrm{Syz}_A^i(A/I) \cong \mathrm{Syz}_A^i(A/J)$ for some $i \geq 0$, where $\mathrm{Syz}_A^i(-)$ denotes the i -th syzygy module in the minimal free resolution. This shows that all the Cohen-Macaulay local rings of finite CM-representation type, i.e., there exist only finitely many non-isomorphic indecomposable maximal Cohen-Macaulay modules, contain only finitely many Ulrich ideals ([5, Theorem 7.8], see also [2, Theorem 5.1]). In particular, by using the techniques from the representation theory of maximal Cohen-Macaulay modules, they succeeded in determining all the Ulrich ideals in Gorenstein local rings of finite CM-representation type of dimension at most 2 ([5, Theorem 9.5], [6, Corollary 5.8], see also [4, Theorem 6.1]). In addition, over a Gorenstein local ring (A, \mathfrak{m}) , if I and J are Ulrich ideals of A with $\mathfrak{m}J \subseteq I \subsetneq J$, then A must be a hypersurface ([5, Corollary 7.6]). Thus the ubiquity of Ulrich ideals reflects the singularities of base rings. Subsequently, the authors of [7] studied the structure of the complex $\mathbf{R}\mathrm{Hom}_A(A/I, A)$ in the derived category of a Cohen-Macaulay local ring A with $d = \dim A$, and proved that A is Gorenstein if and only if A/I is Gorenstein and $\mu_A(I) = d + 1$, if an Ulrich ideal I exists ([5, Corollary 2.6 (b)], [7, Corollary 2.6]), where $\mu_A(-)$ denotes the number of generators.

Motivated by this observation, in this paper we investigate the following question.

Question 1.1. Let A be a Gorenstein ring with $d = \dim A > 0$. How many ideals I of A with $\mathrm{ht}_A I = 1$ which satisfy the ring A/I is Gorenstein and $\mu_A(I) \geq 2$ exist?

2020 *Mathematics Subject Classification.* 13H10, 13A15, 13A02.

Key words and phrases. Gorenstein ring, numerical semigroup ring, a-invariant.

The author was partially supported by JSPS Grant-in-Aid for Young Scientists 20K14299 and JSPS Grant-in-Aid for Scientific Research (C) 23K03058.

Every Ulrich ideal in a one-dimensional Gorenstein local ring satisfies the conditions stated as in Question 1.1. Whereas, based on the preceding researches on Ulrich ideals ([1, 3, 4, 7]), it is rather difficult to make a list of all the Ulrich ideals even for one-dimensional Cohen-Macaulay local rings, especially for numerical semigroup rings; see e.g., [1, Theorem 3.9, Theorem 4.1]. In light of the result that there are only finitely many Ulrich ideals generated by monomials in numerical semigroup rings ([5, Theorem 6.1]), we start our investigation on Question 1.1 by going over graded ideals. Still, it remains unclear the question even if we restrict to graded ideals in numerical semigroup rings, which we will clarify in this paper.

To state our result, let us explain the notation. For a numerical semigroup H , the ring

$$R = k[H] = k[t^h \mid h \in H] \subseteq k[t]$$

is called the *numerical semigroup ring* of H over a field k , where t denotes an indeterminate over k . Note that $R = k[H]$ is a one-dimensional Noetherian graded integral domain; moreover the ring R enjoys a beautiful relation with its corresponding semigroup H . A typical example is that the maximum integer $f(H)$ in the set $\mathbb{N} \setminus H$ coincides with the a -invariant $a(R)$ of the ring $R = k[H]$ ([8, Example (2.1.9)]). Besides, the semigroup H is *symmetric*, i.e., the equality $\#\{n \in H \mid n < c(H)\} = \#(\mathbb{N} \setminus H)$ holds, if and only if its semigroup ring $R = k[H]$ is Gorenstein, where \mathbb{N} denotes the set of non-negative integers, $\#(-)$ is the cardinality of a set, and $c(H) = f(H) + 1$ is the conductor of H . See [10, Proposition 2.21] or [12, Theorem] for the proof of this fact.

With this notation, the main result of this paper is stated as follows.

Theorem 1.2. *Suppose that $R = k[H]$ is a Gorenstein ring. Then the equality*

$$\#\{I \mid I \text{ is a graded ideal of } R \text{ such that } R/I \text{ is Gorenstein and } \mu_R(I) \geq 2\} = c(H)$$

holds.

Let us now explain how this paper is organized. To show Theorem 1.2, we need several auxiliaries which we will prepare in Section 2. We actually provide them in a bit more general setting, not only for numerical semigroup rings. We shall prove Theorem 1.2 in Section 3 starting with the case where $a(R/I) < a(R)$. In Section 4 we finally provide a complete list of non-principal graded ideals I in R whose quotient rings R/I are Gorenstein. As an application of Theorem 1.2, we consider such ideals in the associated graded ring with respect to a certain filtration of ideals. Examples are explored as well.

2. PRELIMINARIES

Let $R = \bigoplus_{n \geq 0} R_n$ be a one-dimensional Noetherian graded integral domain. Throughout this section, we assume $k = R_0$ is a field, and $R_n \neq (0)$ and $R_{n+1} \neq (0)$ for some $n \geq 0$. Let W be the set of non-zero homogeneous elements in R . Note that the localization $W^{-1}R = K[t, t^{-1}]$ of R with respect to W is a *simple* graded ring, i.e., every non-zero homogeneous element is invertible, where t is a homogeneous element of degree 1 which is transcendental over k , and $K = [W^{-1}R]_0$ is a field. There is an exact sequence

$$0 \rightarrow R \rightarrow K[t, t^{-1}] \rightarrow H_{\mathfrak{m}}^1(R) \rightarrow 0$$

of graded R -modules, where \mathfrak{m} denotes the graded maximal ideal of R and $H_{\mathfrak{m}}^1(R)$ is the 1st graded local cohomology module of R with respect to \mathfrak{m} . As $R_0 = k$ and $[H_{\mathfrak{m}}^1(R)]_0$ is a finite-dimensional k -vector space (remember that $H_{\mathfrak{m}}^1(R)$ is an Artinian R -module), the field extension

K/k is finite. Hence $k = K$, if k is an algebraically closed field. Let \bar{R} be the integral closure of R in its quotient field $Q(R)$.

Recall that an integral domain A is called $N-1$ if the integral closure of A in $Q(A)$ is a finite A -module; and $N-2$ if, for any finite field extension L of $Q(A)$, the integral closure of A in L is a finite A -module. We say that a Noetherian ring B is *Nagata* if B/\mathfrak{p} is $N-2$ for every $\mathfrak{p} \in \text{Spec } B$ ([14, (31.A) Definitions]). Note that every field is Nagata, and every finitely generated algebra over a Nagata ring is Nagata ([14, (31.H) Theorem]). Thus every finitely generated algebra over a field which is an integral domain is a Nagata domain, so it is $N-1$. The reader may consult [14, Section 31] for the details.

We begin with the following which was pointed out by S. Goto.

Lemma 2.1. *The equality $\bar{R} = K[t]$ holds in $Q(R)$.*

Proof. Note that \bar{R} is a graded ring and $\bar{R} \subseteq W^{-1}R = K[t, t^{-1}]$; see e.g., [17, page 157]. As the field k is Nagata, so is the finitely generated k -algebra R . Thus \bar{R} is a finite R -module. As $R_n = (0)$ for all $n < 0$ and $R_0 = k$, we see that $[\bar{R}]_n = (0)$ for all $n < 0$, $L = [\bar{R}]_0$ is a field, and $k \subseteq L \subseteq K$. Set $N = \bigoplus_{n>0} [\bar{R}]_n$. Since the local ring \bar{R}_N of \bar{R} at the maximal ideal N is a DVR, the ideal N is principal. We choose a homogeneous element $f \in \bar{R}$ of degree $q > 0$ such that $N = f\bar{R}$. Hence $\bar{R} = L[N] = L[f] \subseteq W^{-1}R = K[t, t^{-1}]$. Besides, because $\bar{R}[f^{-1}] = L[f, f^{-1}]$ is a simple graded ring and $R \subseteq \bar{R}[f^{-1}]$, we have $W^{-1}R \subseteq \bar{R}[f^{-1}] = L[f, f^{-1}]$. Therefore

$$K[t, t^{-1}] = L[f, f^{-1}]$$

so that $K = L$ and $q = 1$. This shows $\bar{R} = L[f] = K[f] = K[t]$, as claimed. \square

For R -submodules X and Y of $Q(R)$, let $X : Y = \{a \in Q(R) \mid aY \subseteq X\}$. If we consider ideals I, J of R , we set $I :_R J = \{a \in R \mid aJ \subseteq I\}$. Hence $I :_R J = (I : J) \cap R$.

Remark 2.2. Let I be a non-zero graded ideal of R . It is straightforward to check that $R : I$ is a graded R -submodule of $K[t, t^{-1}]$ which contains R . In addition, the natural isomorphism $R : I \xrightarrow{\cong} \text{Hom}_R(I, R)$, $\alpha \mapsto (x \mapsto \alpha x)$ is graded. Thus, provided $k = K$, every homogeneous component of $\text{Hom}_R(I, R)$ has dimension, as a k -vector space, at most 1.

For a Cohen-Macaulay graded ring $A = \bigoplus_{n \geq 0} A_n$ such that A_0 is a local ring, we set $a(A) = \max\{n \in \mathbb{Z} \mid [H_{\mathfrak{M}}^d(A)]_n \neq (0)\}$ which is called the *a-invariant* of A ([8, Definition (3.1.4)]). Here, \mathfrak{M} denotes the unique graded maximal ideal of A , $d = \dim A$, and $\{[H_{\mathfrak{M}}^d(A)]_n\}_{n \in \mathbb{Z}}$ is the homogeneous components of the d -th graded local cohomology module $H_{\mathfrak{M}}^d(A)$ of A with respect to \mathfrak{M} . When A admits the graded canonical module K_A , one has $a(A) = -\min\{n \in \mathbb{Z} \mid [K_A]_n \neq (0)\}$.

Let M be a graded R -module and ℓ an integer. Let $M(\ell)$ denote the graded R -module whose underlying R -module is the same as that of the R -module M and the grading is given by $[M(\ell)]_n = M_{\ell+n}$ for all $n \in \mathbb{Z}$. When M is finitely generated, we denote by $\mu_R(M)$ the minimal number of generators of M .

With this notation, we furthermore assume R admits a graded canonical module K_R . Let $(-)^{\vee} = \text{Hom}_R(-, K_R)$ denote the canonical dual functor. We then have the following.

Lemma 2.3. *Suppose that R is a Gorenstein ring. Let I be a graded ideal of R such that R/I is Gorenstein and $\mu_R(I) \geq 2$. Then the following assertions hold true.*

- (1) $[I^\vee]_{-a(R)} \neq (0)$ and $[I^\vee]_{-a(R/I)} \neq (0)$.
- (2) $\min\{n \in \mathbb{Z} \mid [I^\vee]_n \neq (0)\} = \min\{-a(R), -a(R/I)\}$.
- (3) $\mu_R(I^\vee) = 2$.
- (4) If $k = K$, then $a(R) \neq a(R/I)$.
- (5) $I^\vee = Rf + Rg$ for some $f \in [I^\vee]_{-a(R)}$ and $g \in [I^\vee]_{-a(R/I)}$.

Proof. We set $a = a(R)$, $b = a(R/I)$, and $n = \min\{n \in \mathbb{Z} \mid [I^\vee]_n \neq (0)\}$. By taking the functor $(-)^\vee$ to the exact sequence $0 \rightarrow I \rightarrow R \rightarrow R/I \rightarrow 0$, we get the sequence

$$(*) \quad 0 \rightarrow R(a) \rightarrow I^\vee \rightarrow (R/I)(b) \rightarrow 0$$

of graded R -modules, because $K_R \cong R(a)$ and $\text{Ext}_R^1(R/I, K_R) \cong K_{(R/I)} \cong (R/I)(b)$. This shows $[I^\vee]_{-a} \neq (0)$, $[I^\vee]_{-b} \neq (0)$, and $n = \min\{-a, -b\}$. Besides, the exact sequence $(*)$ implies $\mu_R(I^\vee) \leq 2$. As $I^{\vee\vee} \cong I$ and $\mu_R(I) \geq 2$, we get $\mu_R(I^\vee) = 2$. This proves the assertions (1), (2), and (3).

If $k = K$, then all the homogeneous components of $I^\vee \cong \text{Hom}_R(I, R)(a)$, as a k -vector space, have dimension at most 1. Thus $a \neq b$, and the assertion (4) holds.

Recall that \mathfrak{m} is the graded maximal ideal of R . By applying the functor $R/\mathfrak{m} \otimes_R -$ to the sequence $(*)$, we have the exact sequence of the form:

$$(R/\mathfrak{m})(a) \xrightarrow{\xi} I^\vee/\mathfrak{m}I^\vee \xrightarrow{\eta} (R/\mathfrak{m})(b) \rightarrow 0.$$

As $\mu_R(I^\vee) = 2$, the map ξ is injective. We choose $f \in [I^\vee]_{-a}$ and $g \in [I^\vee]_{-b}$ such that $\bar{f} = \xi(1)$ and $\eta(\bar{g}) = 1$, where $\bar{}$ denotes the image in $I^\vee/\mathfrak{m}I^\vee$. Then the images of f, g form a k -basis of $I^\vee/\mathfrak{m}I^\vee$. Hence $I^\vee = Rf + Rg$ by Nakayama's lemma. \square

Remark 2.4. If R is a numerical semigroup ring over a field k , then $k = K$. Whereas, if $k = K$, e.g., k is an algebraically closed field, then the ring R is isomorphic to a semigroup ring of a numerical semigroup ([8, Proposition (2.2.11)]).

3. PROOF OF THEOREM 1.2

We first fix the notation on which all the results in this section are based.

Setup 3.1. Let \mathbb{N} be the set of non-negative integers and $a_1, a_2, \dots, a_\ell \in \mathbb{Z}$ ($\ell \geq 1$) be positive integers such that $\gcd(a_1, a_2, \dots, a_\ell) = 1$. We set

$$H = \langle a_1, a_2, \dots, a_\ell \rangle = \left\{ \sum_{i=1}^{\ell} c_i a_i \mid c_i \in \mathbb{N} \text{ for all } 1 \leq i \leq \ell \right\}$$

and call it the *numerical semigroup* generated by $\{a_i\}_{1 \leq i \leq \ell}$. The reader may consult the book [15] for the fundamental results on numerical semigroups. Let $S = k[t]$ denote the polynomial ring over a field k , and define

$$k[H] = k[t^{a_1}, t^{a_2}, \dots, t^{a_\ell}] \subseteq S$$

which we call the *semigroup ring* of H over k . The ring $R = k[H]$ forms a Noetherian integral domain with $\dim R = 1$ and is a \mathbb{Z} -graded subring of S whose grading $\{R_n\}_{n \in \mathbb{Z}}$ is given by

$$R_n = \begin{cases} kt^n & \text{if } n \in H, \\ (0) & \text{otherwise.} \end{cases}$$

In addition, S is a birational module-finite extension of R , so that $\bar{R} = S$, where \bar{R} denotes the integral closure of R in its quotient field $Q(R)$. Let

$$c(H) = \min\{n \in \mathbb{Z} \mid m \in H \text{ for all } m \in \mathbb{Z} \text{ such that } m \geq n\},$$

and set $f(H) = \max(\mathbb{Z} \setminus H)$ which is called the *Frobenius number* of H . By [8], we get

$$R : S = t^{c(H)}S \text{ and } f(H) = c(H) - 1 = a(R).$$

Note that, for each non-zero ideal I in R , we have $a(R/I) \in H$, whence $a(R/I) \neq a(R)$; see also Lemma 2.3 (4). We set $a = a(R)$ and $c = c(H)$.

Remark 3.2. If R is integrally closed, then $c = 0$ and there is no non-principal ideals in R .

The following plays a key in our argument.

Proposition 3.3. *Suppose that $R = k[H]$ is a Gorenstein ring. Then the equality*

$$\#\{I \mid I \text{ is a graded ideal of } R \text{ such that } a(R/I) < a \text{ and } \mu_R(I) \geq 2\} = \frac{c}{2}$$

holds.

Proof. Let \mathcal{Y}_R be the set of graded ideals I of R such that R/I is Gorenstein, $a(R/I) < a$, and $\mu_R(I) \geq 2$. By Remark 3.2, we may assume $R \neq \bar{R}$. Thus $\mathcal{Y}_R \neq \emptyset$. For each $I \in \mathcal{Y}_R$, since $a(R/I) < a$, we then have $R_m \subseteq I$ for all $m \geq c = a + 1$. By setting $J = R : I$, we see that J is a graded ideal of R and

$$R \subseteq J \subseteq R : \mathfrak{c} = R : (R : \bar{R}) = \bar{R}$$

where the second inclusion follows from $\mathfrak{c} \subseteq I$ and the last equality holds by [11, Bemerkung 2.5] (remember that R is a Gorenstein ring). This shows $J = (1, t^m)$ for some $m \in \mathbb{N} \setminus H$. So we can consider the map

$$\Phi : \mathcal{Y}_R \rightarrow \mathbb{N} \setminus H$$

defined by $\Phi(I) = m$ for each $I \in \mathcal{Y}_R$, where $R : I = (1, t^m)$.

Conversely, for each $m \in \mathbb{N} \setminus H$, we set $J = (1, t^m)$. Then $R \subsetneq J \subseteq \bar{R} = k[t]$. By setting $I = R : J$, we have

$$\mathfrak{c} = R : \bar{R} \subseteq R : J = I \subsetneq R$$

which yield that $a(R/I) < a$, $\mu_R(I) \geq 2$, and the ring R/I is Gorenstein. Indeed, since $t^c \bar{R} = \mathfrak{c} \subseteq I$ and $a(R/I) \in H$, we have $a(R/I) < a$. The K_R -dual $(-)^{\vee}$ of the exact sequence $0 \rightarrow I \rightarrow R \rightarrow R/I \rightarrow 0$ induces the sequence

$$0 \rightarrow R(a) \xrightarrow{\varphi} I^{\vee} \rightarrow \text{Ext}_R^1(R/I, K_R) \rightarrow 0$$

of graded R -modules, because $K_R \cong R(a)$. Let $f = \varphi(1)$. Then $f \in [I^{\vee}]_{-a}$ forms a part of a minimal basis of I^{\vee} . As $I^{\vee} \cong J$ and $\mu_R(J) = 2$, we get $\mu_R(I) \geq 2$; while the R -module $\text{Ext}_R^1(R/I, K_R)$ is cyclic, so that R/I is a Gorenstein ring, because $\text{Ext}_R^1(R/I, K_R) \cong K_{(R/I)}$ is the canonical module of R/I . Here, the proof of above especially shows that if $a(R/I) < a$ then R/I is Gorenstein. Hence, we define the map

$$\Psi : \mathbb{N} \setminus H \rightarrow \mathcal{Y}_R, \quad m \mapsto R : (1, t^m)$$

and it is straightforward to check the composite maps $\Phi \circ \Psi$ and $\Psi \circ \Phi$ are identity. In particular, the map Φ is bijective. Therefore

$$\#\{I \mid I \text{ is a graded ideal of } R, a(R/I) < a, \text{ and } \mu_R(I) \geq 2\} = \#\mathcal{B}_R = \#(\mathbb{N} \setminus H) = \frac{c}{2}$$

where the last equality follows from the fact that H is symmetric, i.e., R is Gorenstein. This completes the proof. \square

We are ready to prove Theorem 1.2.

Proof of Theorem 1.2. Let \mathcal{X}_R be the set of graded ideals I of R such that R/I is Gorenstein and $\mu_R(I) \geq 2$. By Remark 3.2, we may assume $R \neq \bar{R}$. So $\mathcal{X}_R \neq \emptyset$. By Proposition 3.3, it suffices to show that the number of ideals $I \in \mathcal{X}_R$ with $a(R/I) > a$ is a half of the conductor c of H .

For each $I \in \mathcal{X}_R$, we have $\mu_R(I^\vee) = 2$, so we can write

$$I^\vee = Rf + Rg \quad \text{for some } f \in [I^\vee]_{-a} \text{ and } g \in [I^\vee]_{-a(R/I)}.$$

Set $b = a(R/I)$. Since $(0) :_R g = (0)$, we get the exact sequence

$$0 \rightarrow R(b) \xrightarrow{\xi} I^\vee \rightarrow C \rightarrow 0$$

of graded R -modules, where $\xi(1) = g$ and $C = \text{Coker } \xi$. We consider the graded ideal $J = (0) :_R C$ of R . As C has a finite length and $C \cong I^\vee/Rg \neq (0)$, we get $(0) \neq J \subsetneq R$. Besides, we have the isomorphism

$$C \cong R\bar{f} \cong (R/J)(a)$$

as a graded R -module, where $\bar{*}$ denotes the image in I^\vee/Rg . Hence we obtain the sequence

$$0 \rightarrow R(b) \rightarrow I^\vee \rightarrow (R/J)(a) \rightarrow 0$$

of graded R -modules. By applying the functor $(-)^{\vee}$ to the above sequence, we have

$$0 \rightarrow I \rightarrow R(a-b) \rightarrow \text{Ext}_R^1(R/J, K_R)(-a) \rightarrow 0$$

because $I^{\vee\vee} \cong I$ and $R(b)^{\vee} = \text{Hom}_R(R(b), K_R) \cong \text{Hom}_R(R(b), R(a)) \cong R(a-b)$. In particular, $K_{(R/J)} \cong \text{Ext}_R^1(R/J, K_R)$ is cyclic; hence R/J is Gorenstein. By letting $\alpha = a(R/J)$, we have $K_{(R/J)} \cong (R/J)(\alpha)$. Therefore, by changing the shift by $b-a$, we get the exact sequence

$$0 \rightarrow I(b-a) \xrightarrow{\psi} R \rightarrow (R/J)(\alpha-a+b-a) \rightarrow 0$$

of graded R -modules. The degree 0 part of the following isomorphism

$$R/\text{Im } \psi \cong (R/J)(\alpha-2a+b)$$

induces $\alpha-2a+b=0$; while $I(b-a) \cong \text{Im } \psi \cong J$. Hence, $a(R/J) = \alpha = 2a-b$ and $I \cong J(a-b)$ as a graded R -module. In particular, $\mu_R(J) \geq 2$. Thus $J \in \mathcal{X}_R$.

Let W be the set of non-zero homogeneous elements in R . Consider the simple graded ring $W^{-1}R = k[t, t^{-1}]$, where t is a homogeneous element of degree 1 which is transcendental over k . Hence we have the commutative diagram below:

$$\begin{array}{ccc} k[t, t^{-1}](a-b) = W^{-1}(J(a-b)) & \xrightarrow{\cong} & W^{-1}I = k[t, t^{-1}] \\ \uparrow & & \uparrow \\ J(a-b) & \xrightarrow{\cong} & I = t^{b-a}J \end{array}$$

Note that the induced isomorphism $k[t, t^{-1}](a - b) \xrightarrow{\cong} k[t, t^{-1}]$ is given by the homothety of homogeneous element of degree $b - a$. Therefore $I = t^{b-a}J$.

To sum up this argument, for each $I \in \mathcal{X}_R$, there exists a graded ideal $J \in \mathcal{X}_R$ satisfying

$$a(R/J) = 2a - a(R/I) \quad \text{and} \quad I = t^{a(R/I)-a}J.$$

This shows, if $a(R/I) > a$ (resp. $a(R/I) < a$), then $a(R/J) < a$ (resp. $a(R/J) > a$). So, there is a one-to-one correspondence between the set of ideals $I \in \mathcal{X}_R$ with $a(R/I) > a$, and the set of ideals $J \in \mathcal{X}_R$ with $a(R/J) < a$. Finally we conclude that

$$\#\mathcal{X}_R = \#\{I \in \mathcal{X}_R \mid a(R/I) > a\} + \#\{I \in \mathcal{X}_R \mid a(R/I) < a\} = \frac{c}{2} + \frac{c}{2} = c$$

as desired. \square

4. COROLLARIES AND EXAMPLES

We summarize some consequences of Theorem 1.2. In this section we maintain the notation as in Setup 3.1. Let \mathcal{X}_R be the set of graded ideals I of R such that R/I is Gorenstein and $\mu_R(I) \geq 2$. Recall that $a = a(R)$ and $c = c(H)$.

The direct consequence of the proof of Theorem 1.2 is stated as follows, which is useful to compute concrete examples.

Corollary 4.1. *Suppose that $R = k[H]$ is a Gorenstein ring. For each $I \in \mathcal{X}_R$, we set $J = t^{a-a(R/I)}I$. Then the following assertions hold true.*

- (1) $J \in \mathcal{X}_R$ and $a(R/J) = 2a - a(R/I)$. Hence, if $a(R/I) < a$ (resp. $a(R/I) > a$), then $a(R/J) > a$ (resp. $a(R/J) < a$).
- (2) $a(R/I) \in H$, $a \neq a(R/I)$, and $a - a(R/I) \in \mathbb{Z} \setminus H$.
- (3) If $a(R/I) < a$, then $a - a(R/I) \in \mathbb{N} \setminus H$.
- (4) If $a(R/I) > a$, then $a(R/I) - a \in \mathbb{N} \setminus H$.

Proof. We already proved the assertion (1) in the proof of Theorem 1.2. Recall that $a(R/I) \in H$ and $a \notin H$. So $a \neq a(R/I)$. As H is symmetric and $a(R/I) \in H$, we see that $a - a(R/I) \in \mathbb{Z} \setminus H$. In particular, if $a(R/I) < a$, then $a - a(R/I) \in \mathbb{N} \setminus H$. On the other hand, we assume $a(R/I) > a$. Since $J \in \mathcal{X}_R$ and $a(R/J) < a$, we conclude that $a(R/I) - a = a - a(R/J) \in \mathbb{N} \setminus H$, as claimed. \square

The next provides a complete list of graded ideals in \mathcal{X}_R .

Corollary 4.2. *Suppose that $R = k[H]$ is a Gorenstein ring. Then the equality*

$$\mathcal{X}_R = \{R :_R t^m, t^m(R :_R t^m) \mid m \in \mathbb{N} \setminus H\}$$

holds. Moreover, for each $m \in \mathbb{N} \setminus H$, one has

$$a(R/R :_R t^m) = a - m \quad \text{and} \quad a(R/t^m(R :_R t^m)) = a + m.$$

Proof. Note that $R : (1, t^m) = R :_R t^m$ for all $m \in \mathbb{N} \setminus H$. By Proposition 3.3, there is a one-to-one correspondence below:

$$\mathbb{N} \setminus H \longleftrightarrow \{I \in \mathcal{X}_R \mid a(R/I) < a\}, \quad m \longmapsto R : (1, t^m).$$

This shows the equality $\{I \in \mathcal{X}_R \mid a(R/I) < a\} = \{R :_R t^m \mid m \in \mathbb{N} \setminus H\}$. Besides, the proof of Theorem 1.2 guarantees that the map

$$\{I \in \mathcal{X}_R \mid a(R/I) < a\} \longleftrightarrow \{I \in \mathcal{X}_R \mid a(R/I) > a\}, \quad I \longmapsto t^{a-a(R/I)}I$$

is bijective. Hence $\{I \in \mathcal{X}_R \mid a(R/I) > a\} = \{t^{a-a(R/R :_R t^m)}(R :_R t^m) \mid m \in \mathbb{N} \setminus H\}$ holds. Since H is symmetric and $c = a + 1$, it is straightforward to check that $a(R/R :_R t^m) = a - m$ for all $m \in \mathbb{N} \setminus H$. Therefore the equality

$$\mathcal{X}_R = \{R :_R t^m, t^m(R :_R t^m) \mid m \in \mathbb{N} \setminus H\}$$

holds. Furthermore, by Corollary 4.1 (1), we have the equalities

$$a(R/t^m(R :_R t^m)) = 2a - a(R/R :_R t^m) = 2a - (a - m) = a + m$$

which complete the proof. \square

The ideals of the forms $R :_R t^m$ and $t^m(R :_R t^m)$ are easy to compute, especially in numerical semigroup rings, and provide numerous examples illustrating Theorem 1.2.

Example 4.3. Let $k[t]$ be the polynomial ring over a field k and $R = k[H]$ the semigroup ring of a numerical semigroup H . Then the following assertions hold.

(1) Let $H = \langle 2, 2\ell + 1 \rangle$ ($\ell \geq 1$). Then $c(H) = 2\ell$ and the equality

$$\mathcal{X}_R = \{(t^2, t^{2\ell+1}), (t^4, t^{2\ell+1}), \dots, (t^{2\ell}, t^{2\ell+1}), (t^{2\ell+1}, t^{4\ell}), (t^{2\ell+1}, t^{4\ell-2}), \dots, (t^{2\ell+1}, t^{2\ell+2})\}$$

holds.

(2) Let $H = \langle 3, 4 \rangle$. Then $c(H) = 6$ and the equality

$$\mathcal{X}_R = \{(t^3, t^4), (t^4, t^6), (t^3, t^8), (t^8, t^9), (t^6, t^8), (t^4, t^9)\}$$

holds.

(3) Let $H = \langle 3, 5 \rangle$. Then $c(H) = 8$ and the equality

$$\mathcal{X}_R = \{(t^3, t^5), (t^5, t^6), (t^3, t^{10}), (t^5, t^9), (t^{10}, t^{12}), (t^9, t^{10}), (t^5, t^{12}), (t^6, t^{10})\}$$

holds.

(4) Let $H = \langle n, n+1, \dots, 2n-2 \rangle$ ($n \geq 4$). Then $c(H) = 2n$ and the equality

$$\begin{aligned} \mathcal{X}_R &= \{(t^n, t^{n+1}), \dots, (t^{2n-2}, t^{2n-2}), (t^{n+1}, t^{n+2}), \dots, (t^{2n-2}, t^{2n})\} \\ &\cup \{(t^n, t^{n+1}), \dots, (t^{n+i-1}, t^{n+i+1}), \dots, (t^{2n-2}, t^{2n-2}) \mid 1 \leq i \leq n-2\} \\ &\cup \{(t^{3n-1}, t^{3n}), \dots, (t^{4n-3}, t^{4n-3}), (t^{2n}, t^{2n+1}), \dots, (t^{3n-3}, t^{3n-1})\} \\ &\cup \{(t^{2n-i-1}, t^{2n-i}), \dots, (t^{2n-2}, t^{2n}), \dots, (t^{3n-i-3}, t^{3n-i-3}) \mid 1 \leq i \leq n-2\} \end{aligned}$$

holds.

Let $H_1 = \langle a_1, a_2, \dots, a_\ell \rangle$ and $H_2 = \langle b_1, b_2, \dots, b_m \rangle$ ($\ell, m \geq 1$) be numerical semigroups. We choose $d_1 \in H_2 \setminus \{b_1, b_2, \dots, b_m\}$ and $d_2 \in H_1 \setminus \{a_1, a_2, \dots, a_\ell\}$ such that $\gcd(d_1, d_2) = 1$. We say that

$$H = \langle d_1 H_1, d_2 H_2 \rangle = \langle d_1 a_1, d_1 a_2, \dots, d_1 a_\ell, d_2 b_1, d_2 b_2, \dots, d_2 b_m \rangle$$

is a *gluing* of H_1 and H_2 with respect to $d_1 \in H_2$ and $d_2 \in H_1$.

Note that every three-generated symmetric numerical semigroup H is obtained by gluing of a two-generated numerical semigroup H_1 and \mathbb{N} ([9, Section 3], [16, Proposition 3]). Let $a, b \in \mathbb{Z}$

be positive integers with $\gcd(a, b) = 1$. We set $H_1 = \langle a, b \rangle$ and assume that H_1 is minimally generated by two-elements. Choose $c \in H_1$ and $d \in \mathbb{N}$ so that c, d satisfy the conditions that $c > 0$, $d > 1$, $c \notin \{a, b\}$, and $\gcd(c, d) = 1$. Hence, $\gcd(da, db, c) = 1$. We consider a gluing $H = \langle dH_1, c\mathbb{N} \rangle$ of H_1 and \mathbb{N} with respect to $d \in \mathbb{N}$ and $c \in H_1$. Let k be a field. We then have the isomorphism

$$k[H] \cong k[X, Y, Z]/(X^b - Y^a, Z^d - X^m Y^n)$$

of k -algebras, where $c = am + bn$ with $m, n \in \mathbb{N}$. Hence, $a(k[H]) = d(ab - a - b) + (d - 1)c$.

Corollary 4.4. *Let H be a three-generated symmetric numerical semigroup. Under the same notation of above, the equality*

$$\#\mathcal{X}_{k[H]} = d(ab - a - b) + (d - 1)c + 1$$

holds.

Example 4.5. Let k be a field and $H = \langle 4, 6, 7 \rangle$. Then $H = \langle 2\langle 2, 3 \rangle, 7\mathbb{N} \rangle$ and

$$R = k[H] \cong k[X, Y, Z]/(X^3 - Y^2, Z^2 - X^2 Y).$$

In particular, $a(R) = 9$ and $\#\mathcal{X}_R = c(H) = 10$. Indeed, we have the equality

$$\begin{aligned} \mathcal{X}_R &= \{(t^4, t^6, t^7), (t^6, t^7, t^8), (t^4, t^7), (t^4, t^6), (t^6, t^7)\} \\ &\cup \{(t^{13}, t^{15}, t^{16}), (t^{11}, t^{12}, t^{13}), (t^7, t^{10}), (t^6, t^8), (t^7, t^8)\}. \end{aligned}$$

For an R -module M , we denote by $[M]$ the isomorphism class of M .

Corollary 4.6. *Suppose that $R = k[H]$ is a Gorenstein ring. Then the equalities*

$$\{[I] \mid I \in \mathcal{X}_R\} = \{[R :_R t^m] \mid m \in \mathbb{N} \setminus H\} \text{ and } \#\{[I] \mid I \in \mathcal{X}_R\} = \frac{c}{2}$$

hold.

Proof. Note that $R :_R t^m \cong t^m(R :_R t^m)$ as an R -module for each $m \in \mathbb{N} \setminus H$. This shows the equality $\{[I] \mid I \in \mathcal{X}_R\} = \{[R :_R t^m] \mid m \in \mathbb{N} \setminus H\}$ and its cardinality is at most a half of c . We now assume $R :_R t^m = t^\ell(R :_R t^{m'})$ for some $m, m' \in \mathbb{N} \setminus H$ and $\ell \in \mathbb{Z}$. Then

$$R : (1, t^m) = t^\ell(R : (1, t^{m'})) = R : (t^{-\ell}(1, t^{m'})) = R : (t^{-\ell}, t^{-\ell+m'}).$$

As R is Gorenstein, we have $(1, t^m) = (t^{-\ell}, t^{-\ell+m'})$ in $k[t, t^{-1}]$. Since $-\ell < -\ell + m'$, we have $\ell = 0$ and $m = -\ell + m' = m'$. Hence the cardinality of $\{[I] \mid I \in \mathcal{X}_R\}$ is the half of c . \square

Remark 4.7. There exists a one-dimensional local Gorenstein numerical semigroup ring A with infinite residue class field (e.g., $\mathbb{Q}[[t^3, t^7]]$, $\mathbb{C}[[t^4, t^5, t^6]]$) admitting infinitely many two-generated Ulrich ideals. Hence $\mathcal{X}_A = \infty$.

When A is a local ring, although the set \mathcal{X}_A is not necessarily finite, there is an associated graded ring G with respect to a filtration of ideals such that \mathcal{X}_G is a finite set.

Let (A, \mathfrak{m}) be a Noetherian local ring with $\dim A = 1$ and $V = \bar{A}$ the integral closure of A in its total ring $Q(A)$ of fractions. Assume that V is a DVR which is a module-finite extension of A and $A/\mathfrak{m} \cong V/\mathfrak{n}$, where $\mathfrak{n} = tV$ ($t \in V$) denotes the maximal ideal of V . Let $\mathfrak{o}(-)$ denote the \mathfrak{n} -adic valuation (or the order function) of V and set

$$v(A) = \{\mathfrak{o}(f) \mid 0 \neq f \in A\}.$$

Then, $H_A = v(A)$ is called the *value semigroup* of A , which is indeed a numerical semigroup. Let $\mathfrak{c} = A : V$ denote the conductor of A . Then $\mathfrak{c} = t^{c(H_A)}V$ and $c(H_A) = \ell_A(V/\mathfrak{c})$. Note that A is Gorenstein if and only if $H_A = v(A)$ is symmetric ([12, Theorem]).

For each $\ell \in \mathbb{Z}$, we set $F_\ell = \mathfrak{n}^\ell \cap A$. Then $\mathcal{F} = \{F_\ell\}_{\ell \in \mathbb{Z}}$ is a filtration of ideals in A . We define

$$G = G(\mathcal{F}) = \bigoplus_{\ell \geq 0} F_\ell / F_{\ell+1} = \bigoplus_{\ell \geq 0} (\mathfrak{n}^\ell \cap A) / (\mathfrak{n}^{\ell+1} \cap A)$$

and call it the *associated graded ring of A with respect to \mathcal{F}* . Note that, for each $\ell \geq 0$, $G_\ell \neq (0)$ if and only if $\ell \in H_A$. This shows $H_A = \{\ell \geq 0 \mid G_\ell \neq (0)\}$ and the isomorphism below:

$$G = \bigoplus_{\ell \geq 0} F_\ell / F_{\ell+1} \cong (A/\mathfrak{m})[H_A].$$

With this notation we have the following.

Corollary 4.8. *Let (A, \mathfrak{m}) be a one-dimensional Gorenstein complete local domain with algebraically closed residue class field. Let $G = G(\mathcal{F})$ be the associated graded ring of A with respect to the filtration $\mathcal{F} = \{\mathfrak{n}^\ell \cap A\}_{\ell \in \mathbb{Z}}$, where \mathfrak{n} denotes the maximal ideal of $V = \bar{A}$. Then the equality*

$$\#\{I \mid I \text{ is a graded ideal of } G \text{ such that } G/I \text{ is Gorenstein and } \mu_G(I) \geq 2\} = c(H_A)$$

holds, where $H_A = v(A)$ denotes the value semigroup of A .

Acknowledgments. The author would like to thank the referee for his/her valuable comments and suggestions.

REFERENCES

- [1] N. ENDO AND S. GOTO, Ulrich ideals in numerical semigroup rings of small multiplicity, *J. Algebra*, **611** (2022), 435–479.
- [2] N. ENDO AND S. GOTO, Reflexive modules over the endomorphism algebras of reflexive trace ideals, *J. Pure and Appl. Algebra*, **228** (2024), 107662.
- [3] N. ENDO, S. GOTO, S.-I. IAI, AND N. MATSUOKA, Ulrich ideals in the ring $k[[t^5, t^{11}]]$, *Internat. J. Algebra Comput.*, **34** (2024), no.3, 351–369.
- [4] S. GOTO, R. ISOBE, AND N. TANIGUCHI, Ulrich ideals and 2-AGL rings, *J. Algebra*, **555** (2020), 96–130.
- [5] S. GOTO, K. OZEKI, R. TAKAHASHI, K.-I. WATANABE, AND K.-I. YOSHIDA, Ulrich ideals and modules, *Math. Proc. Cambridge Philos. Soc.*, **156** (2014), no.1, 137–166.
- [6] S. GOTO, K. OZEKI, R. TAKAHASHI, K.-I. WATANABE, AND K.-I. YOSHIDA, Ulrich ideals and modules over two-dimensional rational singularities, *Nagoya Math. J.*, **221** (2016), no.1, 69–110.
- [7] S. GOTO, R. TAKAHASHI, AND N. TANIGUCHI, Ulrich ideals and almost Gorenstein rings, *Proc. Amer. Math. Soc.*, **144** (2016), 2811–2823.
- [8] S. GOTO AND K.-I. WATANABE, On graded rings I, *J. Math. Soc. Japan*, **30** (1978), no. 2, 179–213.
- [9] J. HERZOG, Generators and relations of abelian semigroups and semigroup rings, *Manuscripta Math.*, **3** (1970), 175–193.
- [10] J. HERZOG AND E. KUNZ, Die Wertehalbgruppe eines lokalen Rings der Dimnension 1, *S.-Ber. Heidelberger Akad. Wiss. II. Abh.*, 1971.
- [11] J. HERZOG AND E. KUNZ, Der kanonische Modul eines-Cohen-Macaulay-Rings, *Lecture Notes in Mathematics*, **238**, Springer-Verlag, 1971.
- [12] E. KUNZ, The value-semigroup of a one-dimensional Gorenstein ring, *Proc. Amer. Math. Soc.*, **25** (1970), no. 4, 748–751.
- [13] J. LIPMAN, Stable ideals and Arf rings, *Amer. J. Math.*, **93** (1971), 649–685.
- [14] H. MATSUMURA, Commutative algebra, second ed., *Mathematics Lecture Note Series*, **56**, Benjamin / Cummings Publishing Co., Inc., Reading, Mass., 1980.

- [15] J. C. ROSALES AND P. A. GARCÍA-SÁNCHEZ, Numerical semigroups. Developments in Mathematics, 20. *Springer, New York*, 2009.
- [16] K.-I. WATANABE, Some examples of one dimensional Gorenstein domains, *Nagoya Math. J.*, **49** (1973), 101–109.
- [17] O. ZARISKI AND P. SAMUEL, Commutative Algebra Volume II, *Springer*, 1960.

SCHOOL OF POLITICAL SCIENCE AND ECONOMICS, MEIJI UNIVERSITY, 1-9-1 EIFUKU, SUGINAMI-KU,
TOKYO 168-8555, JAPAN

Email address: endo@meiji.ac.jp

URL: <https://www.isc.meiji.ac.jp/~endo/>